

Russian Chemical Bulletin 2008 vol.57 N9, pages 1905-1911

Reactions of heteroaromatic chromophores with lanthanide complexes of p-sulfonatothiocalix[4]arene

Fedorenko S., Mustafina A., Amirov R., Zairov R., Gubaidullin A., Konovalova O., Rusakova N., Kost S., Solovieva S., Antipin I., Konovalov A.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

The conditions for the formation of heteroleptic complexes of a lanthanide(iii) ion ($\text{Ln} = \text{La}, \text{Gd},$ and Tb) with p-sulfonatothiocalix[4]arene and a heteroaromatic chromophore in water were found using X-ray diffraction analysis, pH-metry, ^1H NMR and UV-Vis spectroscopy, and nuclear magnetic relaxation. In the resulting complexes, the heteroaromatic chromophore is in the calix[4]arene cavity and the lanthanide ion is coordinated by the electron-donating groups of the upper or lower calix[4]arene rim. Emission spectroscopic studies revealed changed emission properties of TbIII ions in the terbium(iii)-p-sulfonatothiocalix[4]arene-bipy complex. © 2008 Springer Science+Business Media, Inc.

<http://dx.doi.org/10.1007/s11172-008-0257-7>

Keywords

Chromophores, Heteroleptic complexes, Lanthanides, Nitrogen-containing organic compounds, P-sulfonatothiocalix[4]arene